

Inferring PSC Composition in the Arctic from UARS MLS HNO₃ and POAM II Aerosol Extinction Data

Michelle L. Santee

A. Tabazadeh

G.L. Manney

M.D. Fromm

R.M. Bevilacqua

E.J. Jensen

Overview of Arctic PSC Composition Studies


★☆ Focus on 1995/1996 winter ☆★


Data


- ◆ Polar Ozone and Aerosol Measurement (POAM) II measurements of aerosol extinction at $1.06 \mu\text{m}$
 - ❖ typically 12 solar occultations per day at $\sim 65\text{-}68^\circ\text{N}$
 - ❖ PSCs observed almost daily between mid-December 1995 and early March 1996
- ◆ UARS Microwave Limb Sounder (MLS) measurements of gas-phase HNO_3
 - ❖ Full hemispheric coverage on 22 days during 16 November – 27 December, 1995
 - ❖ Full hemispheric coverage on 17 days during 29 January – 3 March, 1996


Approach


- ◆ The observed evolution of the gas-phase HNO_3 is compared against the observed aerosol extinction and the behavior predicted by various models of PSC composition:
 - ❖ NAT: [Hanson and Mauersberger, 1988]
 - ❖ NAD: [Worsnop et al., 1993]
 - ❖ STS: [Tabazadeh et al., 1994]













17 Feb 1996: POAM rev #12411
POAM PSC Category 3 at 18 km, 433 K, 67.4N, 64.0E



31 Jan 1996: POAM rev #12168
POAM PSC Category 3 at 17 km, 421 K, 66.1N, 92.9E



13 Feb 1996: POAM rev #12356
POAM PSC Category 3 at 20 km, 465 K, 67.2N, 15.5E

